

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

E83-10050

AgRISTARS

"Made available under NASA sponsorship
in the interest of early and wide dis-
semination of Earth Resources Survey
Program information and without liability
for any use made thereof."

Early Warning and Crop Condition Assessment

EW-12-04349 ✓

JSC-18561 ✓

CR-169566

A Joint Program for
Agriculture and
Resources Inventory
Surveys Through
Aerospace
Remote Sensing

SEPTEMBER 1982

DIURNAL PATTERNS OF WHEAT SPECTRAL REFLECTANCES AND
THEIR IMPORTANCE IN THE ASSESSMENT OF CANOPY PARAMETERS
FROM REMOTELY SENSED OBSERVATIONS



✓
P. J. PINTER, JR.; R. D. JACKSON;
S. B. IDS0; R. J. REGINATO

(E83-10050) DIURNAL PATTERNS OF WHEAT
SPECTRAL REFLECTANCES AND THEIR IMPORTANCE
IN THE ASSESSMENT OF CANOPY PARAMETERS FROM
REMOTELY SENSED OBSERVATIONS (Department of
Agriculture, Phoenix, Ariz.) 49 p

N83-14559

Unclas
G3/43 00050

USDA/SRS
U. S. WATER CONSERVATION LABORATORY
4331 E. BROADWAY
PHOENIX, ARIZONA 85040



Lyndon B. Johnson Space Center
Houston, Texas 77058

1. Report No.	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle Diurnal Patterns of Wheat Spectral Reflectances and Their Importance in the Assessment of Canopy Parameters from Remotely Sensed Observations		5. Report Date	
		6. Performing Organization Code	
7. Author(s) P. J. Pinter, Jr. S. B. Idso R. D. Jackson R. J. Reginato		8. Performing Organization Report No.	
		10. Work Unit No.	
9. Performing Organization Name and Address U.S. Water Conservation Laboratory 4331 E. Broadway Phoenix, Arizona 85040		11. Contract or Grant No.	
		13. Type of Report and Period Covered	
12. Sponsoring Agency Name and Address USDA/ARS EW/CCA 1050 Bay Area Blvd., Houston, TX 77058		14. Sponsoring Agency Code	
15. Supplementary Notes			
16. Abstract Spectral reflectances of Produr wheat were measured at 13 different times of the day at Phoenix, Arizona, during April 1979 using a nadir-oriented hand-held 4-band radiometer which had bandpass characteristics similar to those on LANDSAT satellites. Different sun altitude and azimuth angles caused significant diurnal changes in radiant return in both visible and near-IR regions of the spectrum and in several vegetation indices derived from them. The magnitude of these changes were related to different canopy architecture, percent cover and green leaf area conditions. Spectral measurements taken at each time period were well correlated with green leaf area index but the nature of the relationship changed significantly with time of day. Thus, a significant bias in the estimation of GLAI from remotely sensed spectral data could occur if sun angles are not properly accounted for.			
17. Key Words (Suggested by Author(s))		18. Distribution Statement	
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of Pages 49	22. Price*

*For sale by the National Technical Information Service, Springfield, Virginia 22161

Title: DIURNAL PATTERNS OF WHEAT SPECTRAL REFLECTANCES AND THEIR IMPORTANCE IN THE ASSESSMENT OF CANOPY PARAMETERS FROM REMOTELY SENSED OBSERVATIONS

Authors: P. J. Pinter, Jr., Entomologist
R. D. Jackson, Research Physicist
S. B. Idso, Research Physicist
R. J. Reginato, Soil Scientist

Location: U. S. Water Conservation Laboratory, 4331 E. Broadway, Phoenix, Arizona 85040

CRIS No. 5510-20760-001

Publication: Geosciences and Remote Sensing

INTERPRETIVE SUMMARY

Using a hand-held radiometer with bandpass characteristics similar to those on present LANDSAT satellites, spectral reflectance measurements of Produr wheat fields were made at 13 different times of the day at Phoenix, Arizona. The main objective was to determine the effect of changing sun angles on the reflectance properties of canopies in various phenological stages and with different levels of green leaf area index (GLAI). Results indicated that diurnal changes in each of the four LANDSAT wavebands and several indices derived from them are related to canopy architecture, percent cover and vertical distribution of green leaves within the canopy. An analysis using our data showed that substantial errors could be introduced into the estimate of GLAI from spectral observations if the diurnal patterns of reflectance caused by changing sun angles are not properly accounted for.

I. INTRODUCTION

Recent developments in hand-held radiometry [1] have paved the way for the rapid collection of broad waveband spectral data over experimental agricultural plots. Such data are essential to develop and understand the relationships between spectral and agronomic data, such as leaf area index (LAI), biomass and yield. In turn, these relationships should provide needed information to improve the interpretation of satellite data and provide inputs toward the design of future satellite instrumentation and orbital characteristics. Ground-based and aircraft-collected spectral data are not limited to a specific time-of-day acquisition, as are present LANDSAT satellite data. Instead, they are controlled by other constraints, such as those imposed by weather, personnel and equipment availability. Although most researchers limit data collection to several hours before and after solar noon, the time involved for a sequence of measurements over widely spaced fields is often substantial. The bidirectional reflectance properties of plant canopies are complex [2] and not well understood. As a result, spectral data are usually not adjusted for diurnal or seasonal changes in sun angle; a factor which can lead to serious errors or misinterpretation of results.

In an earlier paper [3], we presented spectral data for wheat plots that were hand planted to a high density within rows and approaching a hedgerow type configuration. At the time of our observations all plants were in the same phenological stage. We discussed the complex interactions of sun angles with row orientation, geometry and light reflectance, and concluded that changing sun angles due

to time-of-day were sufficient to obscure apparent spectral differences which might have been attributed to varietal characteristics. Because those results appeared to contradict earlier findings of Lemme and Westin [4], we decided that the effect of changing sun angles warranted more extensive investigation under field conditions. As a consequence, the following report describes a series of hand-held spectral measurements made at 13 times during a single day over differentially water-stressed wheat plots in three different growth stages. Our specific objectives were three-fold: First, to document the diurnal pattern of several widely used vegetation indices derived from several visible and infrared wavebands; second, to determine whether an optimum time-of-day exists for extracting agronomic information from these spectral data; and third, to assess the relative magnitude of errors in scene interpretation which may arise when canopy geometry and changing sun angle configuration, due to time of day, seasonal or latitudinal effects, are not properly accounted for.

II. EXPERIMENTAL APPROACH

Agronomic Methods

In 1978 we initiated a "serial cereal" experiment at Phoenix, Arizona (112 W longitude, 33 N latitude). Three plots of wheat (Triticum durum Desf. var. Produra) were planted sequentially, the first on 1 November 1978, the second on 15 December 1978 and the third on 13 February 1979. Seeds were machine-planted in N-S oriented rows that were 18 cm apart. Each planting was divided into four 12 x 13 m

plots for differential irrigation purposes. Although winter rains prevented the early initiation of differential stress on the first two plantings, by 14 April 1979, the day diurnal spectral measurements were made, significant water stress was evident in at least one plot within each planting date.

The agronomic data were estimated from 5 to 6 twice-weekly samples centered around the 14 April measurement period. Each sample consisted of 6 randomly chosen plants. Green leaf area index (GLAI) was determined using an optically integrating leaf area meter. It is defined here as a dimensionless ratio of the green leaf area (m^2) to the surface area of the soil supporting it (m^2). Fraction brown was the ratio of brown LAI to green LAI. Final yield and plant density were determined for the spectral target areas in each plot at the time of harvest, 1 to 2 months later. Percent green cover was estimated by a dot grid analysis of vertical photographs of the target areas.

Spectral Measurements

Observations were made using an Exotech Model 100 A "LANDSAT Groundtruth Radiometer" with 15° field-of-view lenses. The spectral bandpasses for this radiometer are similar to those of LANDSAT satellites, i.e., MSS4 (0.5 to 0.6 μm), MSS5 (0.6 to 0.7 μm), MSS6 (0.7 to 0.8 μm), and MSS7 (0.8 to 1.1 μm).^{1/} The radiometer was

^{1/} LANDSAT spectral data are expected to follow trends similar to Exotech radiometer data, however, absolute values will undoubtedly be different due to calibration differences and atmospheric attenuation. It should be pointed out that radiance data from the LANDSAT MSS7 is quantitized into 63 different levels instead of the 127 levels used in the other three bands. Thus calculations involving digital data from LANDSAT must be scaled appropriately before any direct comparisons between the two data sets can be made.

nadir oriented and hand-held about 2 m above the soil surface. It was used in conjunction with a microprocessor-based portable data acquisition system that allows simultaneous recording of measurements from each band and subsequent direct output to a computer [5]. Data collection began by recording four readings from a BaSO₄ reflectance plate, six readings from designated target zones within each of three wheat subplots, four readings of the plate, six readings each from three wheat subplots, etc., until all twelve subplots were measured. This sequence, requiring about six minutes to complete, was carried out at thirteen times during the day, beginning at 0753 h and ending at 1758 h MST.

III. RESULTS

Weather and Sun Angles

Meteorological conditions on 14 April were characterized by clear skies and relatively low levels of haze throughout the day. Wind conditions remained calm except during the 1538 h and 1634 h measurement period, when light winds were sufficient to cause the leaves to move slightly. Sun altitude changed markedly during the measurements from 22.5° at 0753 h to a maximum of 65.6° at solar noon, and then dropped to 7.2° at 1758 h MST. Sun azimuth varied from 94° to 276° (Fig. 1).

Agronomic Information

Table 1 presents agronomic data for the experimental wheat plots on the day measurements were made. The plots which have the "1" designation, i.e., 1A, 1B, 1C, and 1D, were phenologically most

advanced with kernel development in the milky ripe stage [6]. The green leaf area had passed its peak and was declining. Plot 1B senesced most rapidly and most of its canopy was brown because of insufficient water. Although the lower canopy elements in plots 1A and 1D were also brown, the upper canopy, flag leaves, heads and awns remained green and effectively obscured the soil and brown leaves from an overhead nadir view. A slight amount of lodging was present in 1A and 1D. A maximum yield of 6010 kg/ha was recorded in 1A.

The "2" plots were planted six weeks later. Plant density was slightly less than in the "1" plots due to different environmental conditions during germination and emergence. By mid-April, these plots had headed and were just past the flowering stage with kernels partially formed. With the exception of 2A, which had some dying of leaves due to limited amount of water, these plots were mostly green, the leaves were oriented vertically and percent green cover was 80-90 percent. Yields from these plots ranged from 2290 to 5680 kg/ha.

Plants in the "3" plots were in stage 7 (second node formed and the next to the last leaf just visible). A few brown leaves were present in these canopies, but inasmuch as green cover ranged from 51 to 73 percent, substantial amounts of soil were viewed by the nadir-looking radiometer. Plots 3C and 3D had yields less than 1000 kg/ha. This was caused primarily by drought stress imposed after the diurnal sequence of radiometric measurements in mid-April. Plot 3B was irrigated one day prior to measurements, and thus the surface soils were wet throughout the study.

Diurnal Changes in Crop Spectral Reflectances

a) Ratio and ND Indices: A number of indices derived from radiometric measurements have been proposed to characterize various agronomic parameters such as plant cover, leaf area index, biomass, and yield [7]. Usually incorporating data from several wavebands, these indices exploit the fact that as the density of the photosynthetically active biomass increases, there is a concomitant increase in near-infrared reflectance and a decrease in visible reflectance. For illustration, we have chosen two such indices that utilize band ratioing techniques which make them relatively independent of illumination intensity and eliminate the irradiance measurements required for the calculation of reflectances. The ratioing of waveband radiances instead of reflectances also circumvents errors in irradiance estimates which are introduced by the non-Lambertian response of BaSO_4 reference panels at low sun angles [8].

Figure 2 shows the ratio of radiances in MSS7 to MSS5 and the Normalized Difference parameter, $\frac{2}{ND} = (\text{MSS7} - \text{MSS5})/(\text{MSS7} + \text{MSS5})$, as a function of time of day for five of our experimental fields which represent a range of canopy conditions. The standard error bars show that these indices change significantly during the day for each vegetated field. They are roughly symmetrical about solar noon and attain their lowest values at that time. We observed the greatest diurnal changes when the green LAI was low and substantial amounts of sunlit soil (fields 2A and 3C) or brown leaves (field 15) were viewed by the

$\frac{2}{ND}$ was originally referred to as the Vegetation Index or VI [9].

radiometer. Note that field 2A has three times the green LAI of 1B and although their spectral data are statistically separable at low sun angles, they are almost identical from 1000 to 1500 h. We found the opposite to be true for fields 1D and 3C. They are statistically indistinguishable before 0800 h or after 1700 h, yet at midday they remain well separated. Such examples underscore the complexities of relating single scene multispectral data to agronomic variables without preliminary knowledge of sun angles and canopy configuration.

b) Visible and Near-IR Light: The diurnal patterns illustrated in Fig. 2 can best be explained by examining the interaction of incident light in the visible (MSS5) and near-IR (MSS7) wavebands with canopy architecture and the changing proportions of soil and green or brown canopy elements. First, consider the behavior of visible light when it strikes a canopy (Fig. 3A). We found that its reflectance^{3/} was relatively uniform throughout the day for canopies with LAI > 3. The same was true in field 3B (not shown in Fig. 2A) where soils were wet from a recent irrigation. On the other hand, the vertical reflectance of visible light from more sparse canopies was very dependent upon the direction of insolation and varied by a factor of 2 with a pronounced maximum at solar noon. At low sun altitudes, with the sun azimuth at right angles to row orientation, only the upper canopy elements were illuminated by direct beam insolation. In most of our fields these elements were predominately green plant tissue (heads, awns, and/or upper

^{3/} The Lambertian response of the BaSO₄ reference panel was unknown and thus not corrected for. Data presented by Kimes and Kirchner [8] indicate that this will result in an overestimation of target reflectance factors at low sun altitudes.

canopy leaves). Consequently, early and late in the day all fields should and did look similar from a reflectance standpoint in visible wavelengths. However, as the sun's altitude increases, and the azimuth angle became more parallel with the rows, progressively more and more sunlight penetrated deeper into the canopy, striking both soil and brown leaf targets which have a much higher reflectance for visible light than do green plant leaves. As a result, scene reflectance was higher at noon in plot 1B because of the illumination of brown canopy elements beneath the partly green flag leaves. The reflectance was higher in 2A and 3C because varying amounts of soil and brown leaves were being illuminated. In the dense canopies, the proportion of illuminated green plant tissue remained essentially constant throughout the day, with the consequence that reflectance remained rather uniform despite changing solar angles.

We found reflectance of near-IR light (MSS7) to behave quite differently, reaching a minimum during midday. The near-IR reflectance from the senescent vegetation in field 1B was distinctly lower, but differences between the other canopies were not as clear-cut as were those we observed in the visible wavebands. There are several explanations for less diurnal change in the near-IR. The first, and most obvious, is that there is less difference in reflectance between bare, dry sunlit soil and green canopy in MSS7 (32 percent vs 37 percent, respectively, at solar noon) than in MSS5 (25 percent vs 2 percent). Thus even though more soil was illuminated at high sun angles, the net effect on reflectance in MSS7 was not as great. A second explanation involves the transmittance properties of green leaves. Although they are relatively opaque to visible light, they are considerably less so

in the longer wavelengths, with near-IR light being transmitted through as many as eight leaf layers before complete attenuation (and hence maximum reflectance) occurs. Thus, at low sun angles, incident light in MSS7 is intercepted by the same number of leaves in low as well as high GLAI situations; and since reflectance is always near maximum, the near-IR does not permit discrimination between the two canopy conditions. However, during midday, maximum near-IR reflectance occurs in fields wherein the intercepted density of leaves in the path of incoming light approaches eight layers, a situation which occurs only under very high green biomass conditions.

c) Other Vegetation Indices: Other types of spectral vegetation indices, such as the Green Vegetation Index (GVI), Soil Brightness Index (SBI), and Perpendicular Vegetation Index (PVI) are linear combinations of two or more wavebands. In order to make these indices more comparable between days, seasons, or latitudes, the radiometric data must be expressed as reflectances or converted by some other normalization procedure so that their absolute values are not proportional to the levels of illumination. If irradiances are not measured directly, as is often the case with satellite and aircraft measurements, they must be estimated by invoking a regular season/time/latitude dependent cosine response function, or by measuring the spectral response of a known relatively unchanging target and then normalizing all unknown targets to that observation. It is important to recognize that these "sun angle corrections" will not remove the diurnal changes we report here, since they do not correct for the

non-Lambertian reflectance of light from a canopy. Indeed, when we converted target radiances to reflectances and calculated GVI and SBI using published empirical coefficients and PVI using a soil background line derived for our soil, we found these three indices also changed significantly with time-of-day.

Correlations With GLAI

Numerous studies have documented correlations between spectral indices and various agronomic characteristics, particularly those related to the canopy's photosynthetically active biomass. Little attention has been given to the influence that sun angles have on these relationships. Our data provided a means to test whether certain times were better than others for monitoring canopy conditions. Using simple linear regressions, we tested whether spectral radiances could be used to predict green leaf area index of all 12 wheat plots at thirteen different times of the day. An examination of the residuals associated with each regression indicated that linear models provided an acceptable characterization of the relation.

a) Individual Wavebands: The coefficients of determination, which indicate the amount of variation in GLAI accounted for by a particular waveband are shown as a function of time of day in Fig. 4A. We found that radiances in both MSS4 and MSS5 accounted for more than 60 percent of the total variability in GLAI regardless of time of day (Fig. 4A). Furthermore, our data show that MSS6 has relatively little information about the photosynthetically active biomass in a canopy, since its coefficient of determination remained lower than 0.20 for all but one time period. Although the 0933 h radiances in the second near-IR

waveband explained more of the variability in GLAI than the visible radiances, its overall performance was lower. It was particularly degraded before that time and after 1400 h, implying that radiance in this waveband cannot discriminate different GLAI at low sun angles.

b) Vegetation Indices: Transforming the spectral data into commonly used vegetation indices improved their ability to predict GLAI (Fig. 4B). Both the ratio of MSS7/MSS5 and ND consistently accounted for more variability in GLAI than did any of the wavebands taken separately. In addition, there does not appear to be any one time-of-day which is best for inferring GLAI. The ratio and ND also remained superior to PVI calculated from either radiances or reflectances. In fact, prior to 0930 or after 1400, PVI did not appear to be very responsive to GLAI at all. We suspect that this was due to the relative importance of MSS7 in determining the value of PVI and the fact that there is a serious reduction in green canopy information content of that band at low sun angles.

c) Temporal Changes in Correlations: Although a relatively good correlation exists between the MSS7/MSS5 ratio and ND, and GLAI throughout the day, the nature of the relation changes significantly with time. To illustrate this point, we show these indices vs GLAI for the solar noon and 1800 h observations in Fig. 5. These two sets of data bound the extremes in sun altitude and azimuth which we observed during our study. Data collected at other time periods were intermediate in nature. Our data set is unique in that it encompasses plants in different stages of growth and senescence conditions as well as different levels of plant water status, yet the standard error of the estimate of

GLAI remained at approximately 0.5 units regardless of the time of day. This is an acceptable error level since the field measurement of GLAI itself, using an optical planimeter, has approximately the same degree of precision associated with it. The slope of the ratio vs GLAI remained constant throughout the day while the intercept approached a maximum at solar noon. As a consequence, a predictive equation based on empirical data collected for one set of sun angle will not be able to predict GLAI well unless the spectral data are taken under similar sun angle conditions and unless row configuration remains approximately the same.

Both the slope and intercept of ND vs GLAI changed with time but the relationships appeared to pivot about the higher LAI values. This implies that under relatively high LAI conditions the value of ND will be similar regardless of time of day.

Potential Errors in GLAI Prediction

The final step in our analysis concerned the determination of how much error could be introduced in the estimation of GLAI from use of these spectral indices if an empirical relationship were derived from measurements at solar noon and then applied to spectral data collected at other times. To perform this error analysis, we first calculated an expected spectral vegetation index for GLAI = 1 thru 4 using the specific relationship between GLAI and the observed ratio $7/5$ and ND at each time. Then we used that expected index value to calculate GLAI using the solar noon relationship. Figure 6 shows the percent error from the true GLAI which occurred when we used spectral data collected at different times of the day. Note that substantial errors are introduced for time periods extending several hours on either side of solar noon, a time period

generally considered "safe" for data collection. Errors introduced at the time of LANDSAT overpass are also significant. Our results show that if the ratio 7/5 data collected at solar noon are used to generate a model for the prediction of GLAI from LANDSAT data (atmospheric attenuation problems aside), it will overpredict GLAI by 25 percent to 35 percent (Fig. 6A). Earlier in the day the problem becomes even more acute. For example, if the 0900 ratio data were used, we would predict a GLAI of almost six for a field where it actually was 4.

In general, the percent errors are greater for low GLAI situations and become more tolerable as GLAI approaches higher values. This was particularly evident when we performed the same type of analysis with ND and found that errors remained less than 10 percent for a $GLAI > 4$ regardless of the time of day. This was evident in Fig. 2B and 3B where values of the normalized difference in plots 1D and 2D remained fairly constant at 0.9 throughout the day. The exact reason for the difference in information content of the ratio of MSS7 to MSS5 and the Normalized Difference is unclear at this point.

IV. CONCLUSIONS AND RECOMMENDATIONS

Although our conclusions may be specific to the bright soil and north-south row orientation used in this experiment, we have shown that the diurnal changes in sun altitude and azimuth angles affect visible and near-IR reflectance measurements of a wheat canopy in a complex manner. This is due to the interaction of the specular component of irradiance with crop geometry, soil, and both green and brown plant tissues. At solar noon, more direct beam sunlight penetrates to the ground in a wheat canopy with N-S oriented rows than at other times of the day, causing

maximum reflectance in visible wavebands and minimum near-IR reflectance. As a consequence, several vegetation indices calculated from radiances in these wavebands attained minimum values at that time. Earlier and later in the day values of these indices were significantly higher which could lead to a serious misinterpretation of canopy conditions if illumination angles and crop geometry are not properly accounted for.

Our findings suggest that the remote detection of low green biomass levels will be enhanced if spectral measurements are made under low sun angle conditions. Also, the ability to differentiate between very high green biomass situations will be facilitated by a high sun altitude and a sun azimuth parallel with row orientation. Finally, seasonal and latitudinal ranges of sun position might impose limitations on the use of remotely sensed spectral data for assessment of dense vegetation conditions in some locations.

Researchers must employ caution when extrapolating empirical relations derived under specific sun angle and row orientation conditions to other latitudes, seasons and times-of-day. Poor correlations between instantaneously acquired satellite spectra and "ground truth" collected over a span of several hours may be a result of rapid diurnal changes we observed in spectral reflectances. Likewise, we believe that some apparent differences or similarities between crops could conceivably result from a systematic bias in spectral data caused by a regular measurement sequence in which one target was consistently viewed under a different set of sun angles.

Much information is contained in the diurnal course of canopy spectral reflectance and it should be more thoroughly investigated

to exploit its potential. The remote sensing community has extracted a great deal of information from our present sunsynchronous LANDSATs. However, we may find that seasonal and latitudinal variation in the sun's relative position could place an upper limit on the amount of useful agricultural information which can be extracted from those vehicles. It is likely that the agricultural community would derive more benefit from geostationary satellite platforms capable of monitoring MSS or Thematic Mapper wavebands on a more continuous basis.

The irradiance levels and target radiance data for each time period are available in tabular form as an appendix in the AgRISTAR'S version of this report.

REFERENCES

- [1] Cipra, J. E., M. F. Baumgardner, E. R. Stoner, and R. B. MacDonald, "Measuring radiance characteristics of soil with a field spectroradiometer, Soil Sci. Soc. Amer. Proc., 35:1014-1017, 1971; Exotech, Inc., LANDSAT (ERTS) groundtruth radiometer model 100-A, Technical description, Exotech, Inc., 1200 Quince Orchard Blvd., Gaithersburg, MD, 1975; Pearson, R. D. and L. D. Miller, Remote mapping of standing crop biomass for estimation of the productivity of the shortgrass prairie, Pawnee National Grassland, Colorado, Proc. 8th Intern. Symp. Remote Sens. Environ., p. 1355-1379, 1972; Weber, F. P., "Development and field test of an ERTS-matched four-channel spectrometer, Presented at Fourth Annual Earth Resources Program Review, pp 120-121, Jan 17-21, 1972; Robinson, B. F., M. E. Bauer, V. P. DeWitt, L. F. Silva, and V. C. Vanderbilt, "A multi-band radiometer for field research," Society of Photo-optical Instrumentation Engineers, Measurements of Optical Radiations 196:8-15, 1979; Jackson, R. D., P. J. Pinter, Jr., R. J. Reginato, and S. B. Idso, "Hand-held radiometry," U. S. Dept. Agric., SEA, ARM-W-19, 66 pp, 1980; Tucker, C. J., W. H. Jones, W. A. Kley, and G. J. Sundstrom, "A three-band hand-held radiometer for field use", Science 211:281-283, 1981.
- [2] Breece, H. T. and R. A. Holmes, "Bidirectional scattering characteristics of healthy green soybean and corn leaves in vivo," Applied Optics, 10: 119-127, 1971; Suits, G. H., "The calculation of the directional reflectance of a vegetative canopy," Remote Sensing Environ., 2:117-125, 1972; Duggin, M. J., "Likely effects of solar elevation on the quantification of changes in vegetation

with maturity using sequential LANDSAT imagery," Applied Optics, 16:521-523, 1977; Bonhomme, R., G. Guyot, and J. Riom, "Relations existant entre reflectance d'un couvert vegetal et sa structure, la hauteur du soleil et l'angle de visee - perspectives de modelisation," In Earth Observation from Space and Management of Planetary Resources. J. Plevin, V. Hood, T. D. Guyene and S. Adamy, eds., Toulouse., pp 319-326, 1978; Rao, V. R., E. J. Beach and A. R. Mack, "Bidirectional reflectance of crops and the soil contribution," Remote Sensing of Environment, 8:115-125, 1979; Kollenkark, J. C., V. A. Vanderbilt, C. S. T. Daughtry, and M. E. Bauer, "Influence of solar illumination angle on soybean canopy reflectance", Applied Optics 21:1179-1184, 1982; Kirchner, J. A., D. S. Kimes and J. E. McMurtrey III, "Variation of directional reflectance factors with structural changes of a developing alfalfa canopy", Applied Optics (In press).

- [3] Jackson, R. D., P. J. Pinter, Jr., S. B. Idso, and R. J. Reginato, "Wheat spectral reflectance: Interactions between crop configuration, sun elevation, and azimuth angle," Applied Optics, 18: 3730-3732, 1979.
- [4] Lemme, G. D. and F. C. Westin, "Landsat - simulating radiometer for agricultural remote sensing," Photogrammetric Engineering and Remote Sensing, 45: 99-103, 1979.
- [5] Allen, R. F., R. D. Jackson, and P. J. Pinter, Jr., "To relate LANDSAT data to U. S. agriculture", Agricultural Engineering, 61:12-14, 1980.

- [6] Large, E. C., "Growth stages in cereals. Illustration of the Feekes Scales," Plant Pathology, 3:128-129, 1954.
- [7] Kauth, R. J. and G. S. Thomas, "The tasseled cap - A graphic description of the spectral-temporal development of agricultural crops as seen by LANDSAT," Proc. Symp. Machine Proc. of Remote Sens. Data, LARS, Purdue, IN., 4B:41-50, 1976; Pearson, R. L., L. D. Miller, and C. J. Tucker, "Hand-held spectral radiometer to estimate graminaceous biomass," Applied Optics, 15: 416-418, 1976; Richardson, A. J. and C. L. Wiegand, "Distinguishing vegetation from soil background information," Photogrammetric Engineering and Remote Sensing, 43:1541-1552, 1977; Tucker, C. J., "Red and photographic infrared linear combination for monitoring vegetation," Remote Sensing of Environment, 8:127-150, 1979; Wiegand, C. L., A. J. Richardson, and E. T. Kanemasu, "Leaf area index estimates for wheat from LANDSAT and their implications for evapotranspiration and crop modeling, Agronomy J., 71: 336-342, 1979; Idso, S. B., P. J. Pinter, Jr., R. D. Jackson, and R. J. Reginato, "Estimation of grain yields by remote sensing of crop senescence rates," Remote Sensing of Environment, 9:87-91, 1980; Pinter, P. J., Jr., R. D. Jackson, S. B. Idso, and R. J. Reginato, "Multidate spectral reflectances as predictors of yield in water stressed small grains," International J. of Remote Sensing, 2:43-48, 1980; Tucker, C. J. and B. N. Holben, "Relationship of spectral data to grain yield variation, Photogrammetric Engineering and Remote Sensing, 46:657-666, 1980.

- [8] Kimes, D. S. and J. A. Kirchner, "Irradiance measurement errors due to the assumption of a Lambertian reference panel," Remote Sensing of Environment, (In press).
- [9] Deering, D. W., J. W. Rouse, Jr., R. H. Haas, and J. A. Schell, "Measuring "forage production" of grazing units from LANDSAT MSS data," In Proc. Tenth International Symp. on Remote Sensing of Environment, pp. 1169-1178, 1975.

Table 1. Agronomic characteristics of experimental *Produra* wheat fields at Phoenix, Arizona, on 14 April 1979.

Plot	Stage [6]	Plant Density (m ²)	Height (cm)	Green LAI	Fraction Green	% Green Cover	Total Dry Biomass (kg/ha)	Final Yield (kg/ha)
1A	11.1	117	100	2.8	.60	81	17420	6010
1B	11.1-11.2	120	100	0.5	.40	48	14930	3530
1C	11.1	118	105	1.5	.51	50	15680	4000
1D	11.1	121	102	3.7	.72	86	19690	5630
2A	10.6	95	86	1.4	.67	59	9080	2290
2B	10.6	117	97	3.2	.84	80	11550	3040
2C	10.6	98	97	2.9	.95	89	9650	3650
2D	10.5-10.6	98	96	4.1	.85	88	8220	5680
3A	7	66	46	1.3	.98	51	1210	1550
3B	7	85	46	2.0	.99	71	1760	3240
3C	7	92	45	1.8	.99	73	1830	860
3D	7	85	42	1.6	.96	66	2040	690

LIST OF FIGURES

1. Sun altitude (degrees above the horizon) and azimuth (degrees clockwise from north) angles for Phoenix, Arizona, on 14 April 1979.
2. Diurnal changes in two vegetation indices derived from spectral radiances in MSS5 and MSS7 for five selected Produr wheat fields on 14 April 1979 at Phoenix, Arizona. Vertical bars define ± 1 standard error. Green LAI is shown in parenthesis.
 - 2a. Radiance ratio of MSS 7 to MSS5.
 - 2b. Radiance normalized difference parameter.
3. Reflectance in MSS5 and MSS7 for selected Produr wheat fields as a function of time of day. Vertical bars on figure 3A represent ± 1 SE. They have been omitted on 3B to reduce clutter.
4. Coefficients of determination calculated for the relationship between green LAI of 12 experimental fields and the radiances measured in each waveband (4a), and four indices calculated from them (4b).
5. The relationships between green LAI and two vegetation indices calculated from spectral data in MSS5 and MSS7 for two times of the day when sun angles were at extremes.
6. Percent error in green LAI when the relation between spectral indices and GLAI at solar noon was used with spectral indices observed at other times of the day.




Figure 1

ORIGINAL PAGE IS
OF POOR QUALITY




Figure 2A

ORIGINAL PAGE IS
OF POOR QUALITY




Figure 2B

ORIGINAL PAGE IS
OF POOR QUALITY




FIGURE 3A

ORIGINAL PAGE IS
OF POOR QUALITY




FIGURE 3B

ORIGINAL PAGE IS
OF POOR QUALITY




Figure 4A

ORIGINAL PAGE IS
OF POOR QUALITY




Figure 4B

ORIGINAL PAGE IS
OF POOR QUALITY




Figure 5A

ORIGINAL PAGE IS
OF POOR QUALITY




Figure 5B

ORIGINAL PAGE IS
OF POOR QUALITY




Figure 6A

ORIGINAL PAGE IS
OF POOR QUALITY




Figure 6B

APPENDIX

The following section documents the radiance (Wm^{-2}) measured in each experimental field. The abbreviation "CL" refers to 95% confidence limits ($1.96 \times$ standard error of 6 observations). The measurement sequence was different than the order presented in the tables: it was STANDARD, 1A, 2A, 3A, STANDARD, 1B, 2B, 3B, STANDARD, 1C, 2C, 3C, STANDARD, 1D, 2D, 3D, STANDARD. The 4 lines of irradiance data shown in the tables were calculated from the mean radiance measured over the 0.6×0.6 m painted BaSO_4 reflectance standard at the beginning and end of each set of 3 consecutive plots. Thus the first line of irradiance values can be used to calculate reflectances in the "A" plot; the second for the "B" plots and so forth.

SERIAL CEREAL

SPECTRAL RADIANCES - PRODURA WHEAT

1

PRODURA WHEAT

DIURNAL RUN DAY 79104

ORIGINAL PAGE IS
OF POOR QUALITY

TIME 753 14 APR 79

PLOT	BAND4 (0.5-0.6)		BAND5 (0.6-0.7)		BAND6 (0.7-0.8)		BAND7 (0.8-1.1)	
	RAD	CL.	RAD	CL.	RAD	CL.	RAD	CL.
1A	1.32	0.17	1.51	0.13	10.60	0.85	21.97	2.63
1B	1.63	0.13	2.22	0.16	9.15	0.45	18.48	0.92
1C	1.37	0.34	1.58	0.41	9.29	2.28	17.94	4.61
1D	1.41	0.25	1.47	0.12	12.22	3.13	24.89	5.70
2A	1.54	0.13	1.76	0.21	10.49	0.82	20.66	2.11
2B	1.12	0.27	1.15	0.33	10.14	1.66	20.70	3.01
2C	1.04	0.12	1.09	0.16	10.68	0.64	22.63	1.51
2D	1.02	0.15	0.97	0.18	10.57	1.43	21.93	3.25
3A	2.15	0.36	2.48	0.52	12.49	2.58	23.81	4.82
3B	1.60	0.29	1.50	0.30	11.84	1.98	22.69	3.59
3C	1.54	0.08	1.48	0.10	12.14	0.78	22.88	1.69
3D	1.86	0.24	1.97	0.37	12.79	1.40	24.18	2.88

TIME -----IRRADIANCE FROM-----
-----BARIUM SULFATE STANDARD-----

750	40.68	49.91	40.82	53.88
	41.37	50.68	41.56	54.84
	41.84	51.04	41.70	55.09
	42.75	52.12	42.59	56.21
757				

RADIANCES IN WATTS / M²

OPERATORS -- PJP

INCOMING SOLAR RADIATION - 328.96 WATTS / M²

SUN ALTITUDE ANGLE - 22.5 DEGREES

SUN AZIMUTH ANGLE - 93.9 DEGREES

1 OF DIURNAL RUN - NO RECORDER PROBLEMS

EXCELLENT SKY CONDITIONS - VERY GOOD VISIBILITY

OVERALL DATA QUALITY --- EXCELLENT

ACTUAL DATA FILE - SC50279104

SUN - DIRECT BEAM UNOBSCURED

HAZE - NONE

CLOUDS - NONE

CLOUD COVER (TENTHS) - 0

WINDSPEED - SLIGHT BREEZE

----- DIRECTION - 90

PRECIPITATION - 0

DEW - NONE

IRRIGATION - 3B

DATE OF THIS PRINTOUT 16 JUN 1982 - PJP

US WATER CONSERVATION LAB, USDA, ARS, PHX AZ

SERIAL CEREAL

SPECTRAL RADIANCES - PRODURA WHEAT

2

PRODURA WHEAT

DIURNAL RUN DAY 79104

ORIGINAL PAGE IS
OF POOR QUALITY

TIME 847 14 APR 79

PLOT	BAND4 (0.5-0.6)		BAND5 (0.6-0.7)		BAND6 (0.7-0.8)		BAND7 (0.8-1.1)	
	RAD	CL	RAD	CL	RAD	CL	RAD	CL
1A	2.08	0.31	2.41	0.24	17.25	2.10	36.07	3.82
1B	2.67	0.46	3.77	0.47	14.31	1.62	28.83	1.89
1C	2.04	0.36	2.45	0.45	13.95	3.10	28.21	6.05
1D	1.83	0.23	2.04	0.14	16.97	3.05	35.47	6.14
2A	2.76	0.23	3.53	0.44	16.38	0.87	31.73	2.38
2B	2.12	0.27	2.16	0.28	16.17	1.85	32.95	3.32
2C	1.61	0.18	1.63	0.20	16.77	1.55	35.89	3.19
2D	1.60	0.31	1.55	0.31	17.80	1.48	37.21	3.21
3A	3.82	0.60	5.08	1.16	17.59	2.22	32.03	4.23
3B	2.78	0.51	2.96	0.96	18.08	1.48	34.60	2.67
3C	2.70	0.19	2.82	0.24	18.82	0.99	36.36	2.36
3D	3.25	0.40	3.80	0.63	18.83	1.57	35.55	3.11

TIME -----IRRADIANCE FROM-----
-----BARIUM SULFATE STANDARD-----

844	66.50	80.40	64.39	84.92
	66.58	80.46	63.97	84.24
	66.71	80.50	63.75	83.61
	67.31	81.25	64.30	84.37
851				

RADIANCES IN WATTS / M²

OPERATORS -- PJP

INCOMING SOLAR RADIATION - 537.073 WATTS / M²

SUN ALTITUDE ANGLE - 33.7 DEGREES

SUN AZIMUTH ANGLE - 102.2 DEGREES

2 OF DIURNAL RUN

SO FAR SO GOOD

OVERALL DATA QUALITY --- EXCELLENT

ACTUAL DATA FILE - SC50379104

SUN - DIRECT BEAM UNOBSURED

HAZE - NONE

CLOUDS - NONE

CLOUD COVER (TENTHS) - 0

WINDSPEED - SLIGHT BREEZE

PRECIPITATION - 0

IRRIGATION - 3R

----- DIRECTION - 90

----- DEW - NONE

DATE OF THIS PRINTOUT 16 JUN 1982 - PJP

SERIAL CEREAL

SPECTRAL RADIANCES - PRODURA WHEAT

3

PRODURA WHEAT

DIURNAL RUN DAY 79104

TIME 933 14 APR 79

PLOT	BAND4 (0.5-0.6)		BAND5 (0.6-0.7)		BAND6 (0.7-0.8)		BAND7 (0.8-1.1)	
	RAD	CL	RAD	CL	RAD	CL	RAD	CL
1A	2.82	0.31	2.98	0.18	22.47	2.11	46.50	3.66
1B	3.65	0.29	5.35	0.35	17.42	1.41	34.64	2.96
1C	2.89	0.52	3.49	0.53	17.57	2.87	34.65	5.47
1D	2.52	0.42	2.75	0.35	22.78	4.67	47.17	9.53
2A	3.93	0.40	5.44	0.73	20.21	1.55	39.59	3.29
2B	2.80	0.27	2.90	0.30	21.16	2.55	43.26	4.53
2C	2.17	0.13	2.15	0.13	21.11	0.89	44.89	2.27
2D	2.19	0.27	2.09	0.32	22.71	1.73	47.36	4.31
3A	5.46	0.89	7.30	1.72	22.61	1.95	41.12	4.53
3B	3.17	0.43	3.31	0.64	20.00	1.82	37.94	3.15
3C	3.76	0.36	4.17	0.41	22.66	1.25	42.67	2.45
3D	4.58	0.48	5.83	0.86	22.22	1.01	41.13	2.16

TIME -----IRRADIANCE FROM-----
-----BARIUM SULFATE STANDARD-----

929	86.12	103.22	81.49	107.58
	85.69	103.24	80.95	107.04
	86.23	103.81	81.51	107.69
	87.30	104.94	82.83	109.37
937				

RADIANCES IN WATTS / M²

OPERATORS --- PJP

INCOMING SOLAR RADIATION - 670.705 WATTS / M²

SUN ALTITUDE ANGLE - 42.9 DEGREES

SUN AZIMUTH ANGLE - 110.7 DEGREES

3 OF DIURNAL RUN

SO FAR SO GOOD

OVERALL DATA QUALITY --- EXCELLENT

ACTUAL DATA FILE - SC50479104

SUN - DIRECT BEAM UNOBSURED

HAZE - NONE

CLOUDS - NONE

CLOUD COVER (TENTHS) - 0

WINDSPEED - SLIGHT BREEZE

PRECIPITATION - 0

IRRIGATION - 3B

----- DIRECTION - 90

DEW - NONE

DATE OF THIS PRINTOUT 16 JUN 1982 - PJP

US WATER CONSERVATION LAB, USDA, ARS, PHX AZ

SERIAL CEREAL

SPECTRAL RADIANCES - PRODURA WHEAT

PRODURA WHEAT

DIURNAL RUN DAY 79104

TIME 1018 14 APR 79

ORIGINAL PAGE IS
OF POOR QUALITY

PLOT	BAND4 (0.5-0.6)		BAND5 (0.6-0.7)		BAND6 (0.7-0.8)		BAND7 (0.8-1.1)	
	RAD	CL	RAD	CL	RAD	CL	RAD	CL
1A	3.34	0.21	3.45	0.20	26.41	1.79	54.69	2.92
1B	4.56	0.50	6.82	0.68	20.32	1.84	39.74	2.71
1C	3.59	0.42	4.36	0.52	19.86	2.92	38.79	6.80
1D	2.72	0.21	2.96	0.28	24.27	3.68	51.26	7.72
2A	5.20	0.50	7.48	0.95	24.01	1.20	46.40	2.80
2B	3.53	0.60	3.83	0.70	24.30	3.03	49.57	5.78
2C	2.82	0.15	2.96	0.14	23.99	2.44	51.05	5.43
2D	2.81	0.37	2.71	0.46	26.59	2.28	55.13	5.02
3A	7.21	1.42	10.31	2.81	26.31	0.94	47.54	3.34
3B	3.99	0.46	4.33	0.78	22.82	1.62	43.49	3.36
3C	5.35	0.80	6.40	0.65	26.62	1.27	49.80	2.53
3D	6.12	0.63	8.12	1.26	26.08	0.96	47.89	1.80

TIME -----IRRADIANCE FROM-----
-----BARIUM SULFATE STANDARD-----

1015	104.62	128.89	101.72	129.92
	105.12	126.29	102.30	130.66
	104.97	126.22	98.76	133.43
	105.84	127.25	99.64	134.67
1022				

RADIANCES IN WATTS / M^2

OPERATORS -- PJP

INCOMING SOLAR RADIATION - 796.327 WATTS / M^2

SUN ALTITUDE ANGLE - 51.3 DEGREES

SUN AZIMUTH ANGLE - 121.4 DEGREES

4 OF DIURNAL RUN

SO FAR SO GOOD

OVERALL DATA QUALITY --- EXCELLENT

ACTUAL DATA FILE - SC50579104

SUN - DIRECT BEAM UNOBSURED

HAZE - NONE

CLOUDS - NONE

CLOUD COVER (TENTHS) - 0

WINDSPEED - SLIGHT BREEZE

PRECIPITATION - 0

IRRIGATION - 3B

----- DIRECTION - 90

DEW - NONE

DATE OF THIS PRINTOUT 16 JUN 1982 - PJP

US WATER CONSERVATION LAB, USDA, ARS, PHX AZ

SERIAL CEREAL

SPECTRAL RADIANCES - PRODURA WHEAT

5

PRODURA WHEAT

DIURNAL RUN DAY 79104

ORIGINAL PAGE IS
OF POOR QUALITY

TIME 1104 14 APR 79

PLOT	BAND4 (0.5-0.6)		BAND5 (0.6-0.7)		BAND6 (0.7-0.8)		BAND7 (0.8-1.1)	
	RAD	CL	RAD	CL	RAD	CL	RAD	CL
1A	3.75	0.24	4.21	0.15	28.75	2.28	60.29	4.15
1B	5.34	0.71	8.47	1.24	22.29	1.77	44.78	3.81
1C	4.54	0.76	5.83	0.69	22.95	3.69	44.61	6.87
1D	3.18	0.18	3.43	0.34	25.50	4.37	56.50	6.20
2A	6.55	0.71	9.84	1.32	26.98	0.92	50.97	1.87
2B	4.28	0.48	4.99	0.53	26.41	2.13	54.42	4.41
2C	3.68	0.38	3.97	0.58	28.35	1.53	59.26	3.23
2D	3.21	0.36	3.29	0.42	29.51	1.84	61.90	3.77
3A	8.27	1.69	11.36	2.58	28.52	1.35	50.94	2.20
3B	4.57	0.51	5.11	0.80	24.86	1.36	47.05	2.41
3C	6.27	0.53	7.74	0.92	28.91	1.08	53.60	2.12
3D	7.96	0.73	10.89	1.58	29.78	0.76	53.47	1.47

TIME -----IRRADIANCE FROM-----
-----BARIUM SULFATE STANDARD-----

1101	119.14	142.96	111.81	147.00
	119.05	142.83	111.71	147.16
	119.45	143.11	112.09	147.63
	119.35	143.10	111.71	147.50
1107				

RADIANCES IN WATTS / M²

OPERATORS -- PJP

INCOMING SOLAR RADIATION - 858.407 WATTS / M²

SUN ALTITUDE ANGLE - 58.8 DEGREES

SUN AZIMUTH ANGLE - 136.4 DEGREES

5 OF DIURNAL RUN

SKY CONDITIONS REMAIN EXCELLENT - SO FAR SO GOOD

OVERALL DATA QUALITY *** EXCELLENT

ACTUAL DATA FILE - SC50679104

SUN - DIRECT BEAM UNOBSERVED

HAZE - NONE

CLOUDS - NONE

CLOUD COVER (TENTHS) - 0

WINDSPEED - SLIGHT BREEZE

PRECIPITATION - 0

IRRIGATION - 38

****- DIRECTION - 90

****- DEW - NONE

DATE OF THIS PRINTOUT 16 JUN 1982 - PJP

US WATER CONSERVATION LAB, USDA, ARS, PHX AZ

SERIAL CEREAL

SPECTRAL RADIANCES - PRODURA WHEAT

6

PRODURA WHEAT

DIURNAL RUN DAY 79104

ORIGINAL PAGE 13
OF POOR QUALITY

TIME 1153 14 APR 79

PLOT	BAND4 (0.5-0.6)		BAND5 (0.6-0.7)		BAND6 (0.7-0.8)		BAND7 (0.8-1.1)	
	RAD	CL	RAD	CL	RAD	CL	RAD	CL
1A	4.16	0.14	4.54	0.20	29.96	1.03	61.75	1.90
1B	6.30	0.46	10.00	0.68	23.63	1.32	45.32	3.46
1C	5.33	0.53	7.07	0.69	24.03	2.39	45.52	5.61
1D	3.27	0.06	3.76	0.43	27.24	3.40	56.92	7.21
2A	7.85	0.96	12.14	1.64	29.20	1.20	55.20	1.90
2B	5.04	0.62	5.92	0.86	29.32	1.51	53.48	3.54
2C	4.07	0.54	4.60	0.82	29.47	1.29	61.26	2.47
2D	3.53	0.17	3.59	0.30	29.68	1.59	60.71	3.55
3A	8.84	1.57	12.30	2.25	30.67	0.80	54.56	1.56
3B	5.34	0.37	6.17	0.69	26.41	0.90	49.52	2.08
3C	6.80	0.45	9.08	1.07	29.76	1.17	55.52	2.04
3D	9.19	0.45	12.71	0.97	32.08	0.51	56.65	1.28

TIME -----IRRADIANCE FROM-----
-----BARIUM SULFATE STANDARD-----

1150	130.32	155.51	121.02	158.38
	129.84	154.90	120.20	157.59
	129.80	154.13	119.68	156.53
	130.61	155.36	120.80	158.33
1157				

RADIANCES IN WATTS / M^2

OPERATORS --- PJP

INCOMING SOLAR RADIATION - 924.492 WATTS / M^2

SUN ALTITUDE ANGLE - 64.3 DEGREES

SUN AZIMUTH ANGLE - 159.3 DEGREES

6 OF DIURNAL RUN

SKY CONDITIONS REMAIN EXCELLENT - SO FAR SO GOOD
OVERALL DATA QUALITY --- EXCELLENT

ACTUAL DATA FILE - SC50779104

SUN - DIRECT BEAM UNOBSURED

HAZE - NONE

CLOUDS - NONE

CLOUD COVER (TENTHS) - 0

WINDSPEED - SLIGHT BREEZE

PRECIPITATION - 0

IRRIGATION - 3R

----- DIRECTION - 90

DEW - NONE

DATE OF THIS PRINTOUT 16 JUN 1982 - PJP

US WATER CONSERVATION LAB, USDA, ARS, PHX AZ

SERIAL CEREAL

ORIGINAL PAGE IS
OF POOR QUALITY

SPECTRAL RADIANCES - PRODURA WHEAT

7

PRODURA WHEAT

DIURNAL RUN DAY 79104

TIME 1234 14 APR 79

PLOT	BAND4		BAND5		BAND6		BAND7	
	(0.5-0.6)		(0.6-0.7)		(0.7-0.8)		(0.8-1.1)	
	RAD	CL	RAD	CL	RAD	CL	RAD	CL
1A	4.34	0.13	4.77	0.24	30.57	1.83	62.94	3.59
1B	6.71	0.67	10.90	1.32	23.87	1.55	44.89	4.28
1C	5.51	0.71	7.63	0.80	22.47	3.41	42.69	7.40
1D	3.70	0.37	4.28	0.71	27.55	2.64	58.29	5.82
2A	8.14	0.62	12.73	1.04	29.29	0.70	54.38	2.22
2B	5.52	0.70	6.75	1.21	30.31	1.33	60.28	3.12
2C	4.70	0.53	5.57	0.79	29.98	1.29	61.09	2.39
2D	3.72	0.35	3.89	0.38	30.58	1.50	63.36	3.44
3A	9.80	2.00	14.19	3.83	31.17	0.99	54.45	1.94
3B	5.36	0.32	6.08	0.68	27.50	1.66	51.45	3.67
3C	6.98	0.61	9.55	1.69	30.32	0.66	55.95	1.68
3D	9.45	0.60	13.21	1.48	32.51	0.23	57.54	0.81

TIME -----IRRADIANCE FROM-----
-----BARIUM SULFATE STANDARD-----

1230	132.29	157.64	122.27	160.12
	131.86	156.98	121.59	159.27
	131.95	157.22	121.63	159.82
	132.04	157.19	121.79	160.04
1238				

RADIANCES IN WATTS / M²

OPERATORS -- PJP

INCOMING SOLAR RADIATION - 928.189 WATTS / M²

SUN ALTITUDE ANGLE - 65.6 DEGREES

SUN AZIMUTH ANGLE - 183.3 DEGREES

7 OF DIURNAL RUN - NEW RECORDER BAT PACK

SKY CONDITIONS REMAIN EXCELLENT - SO FAR SO GOOD

OVERALL DATA QUALITY --- EXCELLENT

ACTUAL DATA FILE - SC50879104

SUN - DIRECT BEAM UNOBSERVED

HAZE - NONE

CLOUDS - NONE

CLOUD COVER (TENTHS) - 0

WINDSPEED - SLIGHT BREEZE

PRECIPITATION - 0

IRRIGATION - 3B

----- DIRECTION - 90

DEW - NONE

DATE OF THIS PRINTOUT 16 JUN 1982 - PJP

US WATER CONSERVATION LAB, USDA, ARS, PHX AZ

SERIAL CEREAL

SPECTRAL RADIANCES - PRODURA WHEAT

8

PRODURA WHEAT

DIURNAL RUN DAY 79104

ORIGINAL PAGE 13
OF POOR QUALITY

TIME 1331 14 APR 79

PLOT	BAND4 (0.5-0.6)		BAND5 (0.6-0.7)		BAND6 (0.7-0.8)		BAND7 (0.8-1.1)	
	RAD	CL	RAD	CL	RAD	CL	RAD	CL
1A	4.00	0.18	4.39	0.23	30.15	1.14	63.33	2.99
1B	5.84	0.66	9.31	1.07	22.75	1.90	44.47	4.59
1C	4.87	0.53	6.43	0.60	22.80	2.61	43.38	6.17
1D	3.16	0.18	3.51	0.35	26.93	1.63	57.59	4.06
2A	7.22	0.68	11.15	1.34	27.47	0.86	51.69	1.61
2B	4.77	0.44	5.93	0.54	28.29	0.55	56.98	2.62
2C	3.99	0.64	4.49	0.90	28.32	1.96	58.25	3.54
2D	3.25	0.18	3.25	0.29	29.79	1.31	62.52	2.75
3A	9.02	1.41	12.73	2.65	30.27	0.84	53.79	2.48
3B	5.06	0.28	5.66	0.55	26.93	1.34	50.52	3.08
3C	6.06	0.37	7.89	1.10	29.15	1.49	54.95	3.25
3D	8.18	0.55	11.12	1.29	30.79	0.63	56.10	2.04

TIME -----IRRADIANCE FROM-----
-----BARIUM SULFATE STANDARD-----

1327	126.08	150.09	116.07	152.71
	124.01	147.79	114.49	150.58
	124.06	148.09	114.79	151.53
	124.56	148.54	115.31	152.30
1335				

RADIANCES IN WATTS / M²

OPERATORS -- PJP

INCOMING SOLAR RADIATION - 892.528 WATTS / M²

SUN ALTITUDE ANGLE - 61.7 DEGREES

SUN AZIMUTH ANGLE - 214.2 DEGREES

8 OF DIURNAL RUN

SKY CONDITIONS REMAIN EXCELLENT - SO FAR SO GOOD

OVERALL DATA QUALITY --- EXCELLENT

ACTUAL DATA FILE - SC50979104

SUN - DIRECT BEAM UNOBSURED

HAZE - NONE

CLOUDS - NONE

CLOUD COVER (TENTHS) - 0

WINDSPEED - SLIGHT BREEZE

----- DIRECTION - 90

PRECIPITATION - 0

----- DEW - NONE

IRRIGATION - 38

DATE OF THIS PRINTOUT 16 JUN 1982 - PJP

US WATER CONSERVATION LAB, USDA, ARS, PHX AZ

SERIAL CEREAL

SPECTRAL RADIANCES -- PRODURA WHEAT

9

PRODURA WHEAT

DIURNAL RUN DAY 79104

TIME 1436 14 APR 79

PLOT	BAND4 (0.5-0.6)		BAND5 (0.6-0.7)		BAND6 (0.7-0.8)		BAND7 (0.8-1.1)	
	RAD	CL	RAD	CL	RAD	CL	RAD	CL
1A	3.65	0.22	3.84	0.21	28.18	1.69	57.99	2.57
1B	4.69	0.73	6.97	1.09	20.34	1.81	38.65	3.98
1C	3.56	0.39	4.82	0.79	18.76	2.46	36.39	5.34
1D	2.72	0.24	3.07	0.34	23.33	1.94	49.36	4.81
2A	5.25	0.38	7.77	0.61	23.90	0.74	46.28	1.93
2B	3.50	0.34	3.95	0.36	24.59	1.70	50.19	3.85
2C	3.13	0.39	3.36	0.52	24.49	2.51	50.99	4.95
2D	2.54	0.40	2.57	0.41	26.24	3.28	55.52	7.15
3A	7.28	1.13	10.00	1.67	27.30	1.45	49.50	2.74
3B	4.31	0.24	4.54	0.45	25.49	1.58	48.67	3.01
3C	4.98	0.24	5.89	0.67	26.84	1.50	50.82	3.05
3D	6.36	0.47	8.12	0.90	28.25	1.43	52.51	3.16

TIME -----IRRADIANCE FROM-----
-----BARIUM SULFATE STANDARD-----

1433	107.81	128.73	100.33	132.19
	106.79	127.68	99.56	131.24
	105.67	126.57	98.42	130.01
	105.10	125.75	97.99	129.36
1440				

RADIANCES IN WATTS / M²

OPERATORS -- PJP

INCOMING SOLAR RADIATION - 773.606 WATTS / M²

SUN ALTITUDE ANGLE - 51.9 DEGREES

SUN AZIMUTH ANGLE - 237.7 DEGREES

9 OF DIURNAL RUN

SKY CONDITIONS REMAIN EXCELLENT - SO FAR SO GOOD
OVERALL DATA QUALITY ***- EXCELLENT

ACTUAL DATA FILE - SC51079104

SUN - DIRECT BEAM UNOBSCURED

HAZE - NONE

CLOUDS - NONE

CLOUD COVER (TENTHS) - 0

WINDSPEED - SLIGHT BREEZE

PRECIPITATION - 0

IRRIGATION - 3B

****- DIRECTION - 180

****- DEW - NONE

DATE OF THIS PRINTOUT 16 JUN 1982 - PJP

US WATER CONSERVATION LAB, USDA, ARS, PHX AZ

SERIAL CEREAL

ORIGINAL PAGE IS
OF POOR QUALITY

10

SPECTRAL RADIANCES - PRODURA WHEAT

PRODURA WHEAT

DIURNAL RUN DAY 79104

TIME 1538 14 APR 79

PLOT	BAND4		BAND5		BAND6		BAND7	
	(0.5-0.6)		(0.6-0.7)		(0.7-0.8)		(0.8-1.1)	
	RAD	CL	RAD	CL	RAD	CL	RAD	CL
1A	2.59	0.28	2.88	0.41	21.17	1.58	44.57	3.75
1B	3.34	0.46	4.74	0.59	15.99	2.12	30.97	2.96
1C	2.85	0.28	3.59	0.78	16.92	0.99	32.35	2.77
1D	2.12	0.23	2.27	0.24	17.95	1.39	37.88	2.86
2A	3.35	0.32	4.79	0.41	17.79	1.56	35.87	2.81
2B	2.64	0.42	2.77	0.37	20.37	1.88	41.71	3.31
2C	1.97	0.21	2.05	0.19	17.51	2.06	37.16	3.16
2D	1.54	0.18	1.43	0.20	17.49	1.63	37.68	4.02
3A	5.11	0.76	6.58	1.23	22.82	1.62	42.91	3.73
3B	3.36	0.31	3.32	0.43	21.43	1.40	41.57	2.76
3C	3.45	0.11	3.81	0.26	22.02	1.66	42.71	3.38
3D	4.53	0.32	5.63	0.57	21.92	2.01	41.27	3.69

TIME -----IRRADIANCE FROM-----
-----BARIUM SULFATE STANDARD-----

1535	81.83	98.13	77.32	102.39
	80.31	96.33	75.80	100.46
	79.31	95.30	75.07	99.84
	78.17	94.12	74.14	98.67
1542				

RADIANCES IN WATTS / M^2

OPERATORS -- PJP

INCOMING SOLAR RADIATION - 610.089 WATTS / M^2

SUN ALTITUDE ANGLE - 40.2 DEGREES

SUN AZIMUTH ANGLE - 252.1 DEGREES

10 OF DIURNAL RUN - WIND PICKING UP

SKY CONDITIONS REMAIN EXCELLENT - SO FAR SO GOOD

OVERALL DATA QUALITY ***- EXCELLENT

ACTUAL DATA FILE - SC51179104

SUN - DIRECT BEAM UNOBSCURED

HAZE - NONE

CLOUDS - NONE

CLOUD COVER (TENTHS) - 0

WINDSPEED - LIGHT WIND

PRECIPITATION - 0

IRRIGATION - 3B

***** DIRECTION - 235

***- DEW - NONE

DATE OF THIS PRINTOUT 16 JUN 1982 - PJP

SERIAL CEREAL

SPECTRAL RADIANCES - PRODURA WHEAT

11

PRODURA WHEAT

DIURNAL RUN DAY 79104

TIME 1634

14 APR 79

ORIGINAL PAGE IS
OF POOR QUALITY

PLOT	BAND4 (0.5-0.6)		BAND5 (0.6-0.7)		BAND6 (0.7-0.8)		BAND7 (0.8-1.1)	
	RAD	CL	RAD	CL	RAD	CL	RAD	CL
1A	1.91	0.21	2.14	0.35	15.40	1.63	32.40	3.98
1B	2.34	0.25	3.24	0.41	11.42	1.05	22.59	2.01
1C	1.61	0.23	2.20	0.51	10.03	0.92	20.88	1.67
1D	1.48	0.28	1.60	0.16	12.64	1.85	26.69	2.34
2A	2.31	0.22	3.04	0.38	13.33	0.90	26.69	1.84
2B	1.63	0.16	1.67	0.24	14.39	1.57	29.56	3.31
2C	1.14	0.20	1.12	0.19	10.99	2.00	23.78	3.70
2D	1.21	0.18	1.13	0.18	12.67	1.65	27.18	3.32
3A	3.04	0.43	3.65	0.67	15.99	2.16	30.26	3.86
3B	2.27	0.29	2.22	0.32	15.27	1.25	29.53	2.80
3C	2.24	0.15	2.41	0.15	15.16	1.88	29.11	3.53
3D	2.82	0.25	3.16	0.33	15.88	1.83	30.20	3.63

TIME

-----IRRADIANCE FROM-----
-----BARIUM SULFATE STANDARD-----

1631	55.31	67.00	53.43	70.78
	54.35	65.82	52.58	69.71
	53.52	64.96	51.89	69.03
	53.03	64.49	51.46	68.63
1638				

RADIANCES IN WATTS / M²

OPERATORS -- PJP

INCOMING SOLAR RADIATION - 420.616 WATTS / M²

SUN ALTITUDE ANGLE - 28.8 DEGREES

SUN AZIMUTH ANGLE - 261.6 DEGREES

11 OF DIURNAL RUN - SLIGHTLY CALMER NOW

SKY CONDITIONS REMAIN EXCELLENT - SO FAR SO GOOD

OVERALL DATA QUALITY --- EXCELLENT

ACTUAL DATA FILE - SC51279104

SUN - DIRECT BEAM UNOBSURED

HAZE - NONE

CLOUDS - NONE

CLOUD COVER (TENTHS) - 0

WINDSPEED - LIGHT WIND

PRECIPITATION - 0

IRRIGATION - 3B

----- DIRECTION - 135

----- DEW - NONE

DATE OF THIS PRINTOUT 16 JUN 1982 - PJP

US WATER CONSERVATION LAB, USDA, ARS, PHX AZ

SPECTRAL RADIANCES - PRODURA WHEAT

PRODURA WHEAT

DIURNAL RUN DAY 79104

TIME 1721 14 APR 79

PLOT	BAND4 (0.5-0.6)		BAND5 (0.6-0.7)		BAND6 (0.7-0.8)		BAND7 (0.8-1.1)	
	RAD	CL	RAD	CL	RAD	CL	RAD	CL
1A	1.35	0.27	1.58	0.36	10.85	1.40	22.41	2.93
1B	1.45	0.22	2.13	0.29	7.33	1.23	14.53	1.90
1C	1.13	0.28	1.46	0.42	6.94	1.67	13.74	2.78
1D	0.98	0.17	1.14	0.13	8.57	1.21	18.64	2.79
2A	1.31	0.09	1.61	0.17	8.22	0.61	16.80	1.89
2B	1.05	0.16	1.08	0.20	8.97	1.28	18.61	2.70
2C	0.74	0.12	0.80	0.16	7.15	1.40	15.70	2.87
2D	0.69	0.15	0.69	0.17	7.30	1.21	16.21	3.02
3A	1.72	0.25	1.97	0.32	10.09	1.89	19.79	4.03
3B	1.49	0.17	1.47	0.15	10.45	1.58	20.27	2.63
3C	1.31	0.16	1.32	0.17	9.95	1.11	19.86	2.84
3D	1.57	0.20	1.70	0.20	9.73	1.52	18.93	2.88

TIME -----IRRADIANCE FROM-----
-----BARIUM SULFATE STANDARD-----

1717	33.18	40.93	33.69	44.90
	32.05	39.56	32.58	43.52
	31.38	38.81	31.85	42.82
	30.83	38.12	31.36	42.19

1725

RADIANCES IN WATTS / M²

OPERATORS -- PJP

INCOMING SOLAR RADIATION - 256.328 WATTS / M²

SUN ALTITUDE ANGLE - 19 DEGREES

SUN AZIMUTH ANGLE - 268.5 DEGREES

12 OF DIURNAL RUN - SHADOWS WEST AREAS THIS AND EARLIER RUN

SKY CONDITIONS REMAIN EXCELLENT - SO FAR SO GOOD

OVERALL DATA QUALITY *** EXCELLENT

ACTUAL DATA FILE - SC51379104

SUN - DIRECT BEAM UNOBSCURED

HAZE - NONE

CLOUDS - NONE

CLOUD COVER (TENTHS) - 0

WINDSPEED - NONE

**** DIRECTION - 180

PRECIPITATION - 0

**** DEW - NONE

IRRIGATION - 3B

DATE OF THIS PRINTOUT 16 JUN 1982 - PJP

SERIAL CEREAL

SPECTRAL RADIANCES - PRODURA WHEAT

13

PRODURA WHEAT

DIURNAL RUN DAY 79104

ORIGINAL PAGE IS
OF POOR QUALITY

TIME 1758 14 APR 79

PLOT	BAND4 (0.5-0.6)		BAND5 (0.6-0.7)		BAND6 (0.7-0.8)		BAND7 (0.8-1.1)	
	RAD	CL	RAD	CL	RAD	CL	RAD	CL
1A	0.78	0.10	1.01	0.22	6.59	0.79	14.09	2.07
1B	0.79	0.09	1.10	0.14	4.09	0.59	7.89	0.90
1C	0.59	0.17	0.72	0.26	3.56	1.37	6.81	2.01
1D	0.53	0.09	0.62	0.05	4.56	0.97	9.61	1.96
2A	0.69	0.07	0.82	0.10	4.60	0.45	9.51	1.14
2B	0.42	0.03	0.43	0.08	3.79	0.47	7.96	1.01
2C	0.38	0.10	0.39	0.10	3.62	0.64	7.78	1.20
2D	0.32	0.07	0.31	0.09	3.65	0.69	7.84	1.79
3A	0.83	0.13	0.91	0.16	5.22	1.24	10.50	2.64
3B	0.73	0.06	0.72	0.03	5.24	0.73	10.14	1.14
3C	0.68	0.10	0.68	0.10	5.21	1.03	10.14	2.00
3D	0.74	0.08	0.80	0.11	4.78	0.99	9.58	2.17

TIME -----IRRADIANCE FROM-----
-----BARIUM SULFATE STANDARD-----

1755	16.47	20.62	17.70	23.92
	15.80	19.85	17.13	23.12
	15.26	19.22	16.57	22.34
	14.59	18.37	15.92	21.49
1802				

RADIANCES IN WATTS / M²

OPERATORS -- PJP

INCOMING SOLAR RADIATION - 134.788 WATTS / M²

SUN ALTITUDE ANGLE - 7.2 DEGREES

SUN AZIMUTH ANGLE - 276.3 DEGREES

13 OF DIURNAL RUN - SOME SKINNY SHADOWS FROM POLES ONLY
SKY CONDITIONS REMAIN EXCELLENT - SO FAR SO GOOD
OVERALL DATA QUALITY *** EXCELLENT

ACTUAL DATA FILE - SC51479104

SUN - DIRECT BEAM UNOBSCURED

HAZE - NONE

CLOUDS - NONE

CLOUD COVER (TENTHS) - 0

WINDSPEED - NONE

****- DIRECTION - 180

PRECIPITATION - 0

****- DEW - NONE

IRRIGATION - 3B

DATE OF THIS PRINTOUT 16 JUN 1982 - PJP

US WATER CONSERVATION LAB, USDA, ARS, PHX AZ

U.S. GOVERNMENT PRINTING OFFICE: 1982-289-012123